



Osteopathic Manipulative Treatment (OMT) of the Cervical Spine

Policy Statement

The American Osteopathic Association, in the hopes of advancing the science of osteopathic medicine adopts the following position:

These recommendations are provided for osteopathic educators and physicians making decisions regarding the instruction of cervical spinal manipulation and the care of patients. As such, they cannot substitute for the individual judgment brought to each clinical situation by a patient's physician. Like all reference resources, they reflect the best understanding of the science of medicine at the time of publication, but they should be used with the understanding that continued research is needed.

AMERICAN OSTEOPATHIC ASSOCIATION OSTEOPATHIC MANIPULATIVE TREATMENT OF THE CERVICAL SPINE

Background and Statement of Issue

Treating chronic pain continues to be an important health issue for osteopathic physicians. Chronic pain affects over 100 million Americans over the age of 18 and negatively impacts their quality of life.¹ In addition, it costs \$600 billion a year in healthcare costs and loss of productivity.¹ Back and neck pain are two leading causes of chronic pain and they are amongst the leading causes of people living with disabilities in the United States (U.S.) as well as worldwide. More specifically, back and neck pain are ranked in the top 8 diseases and injuries in the U.S. regarding years lived with disability (YLDs)² and in the top 6 globally.³ Cervical spine manipulation is one option for treating back and neck pain.

Concerns continue to arise regarding the safety of cervical spine manipulation. Specifically, concerns center on the potential development of serious adverse events such as stroke and cervical artery dissection after spinal manipulation. Since spinal manipulation is an option available to osteopathic physicians to incorporate into the care of their patients, it is important to examine these concerns and develop a position on the issue. This paper will present the evidence behind the benefit of cervical spine manipulation, explore the potential harms and make a recommendation about its use.

Benefit

Spinal manipulation has been reviewed in various systematic reviews and meta-analyses over the past three decades. The majority of the studies conducted on spinal manipulation focus on low back pain for which the evidence has shown spinal manipulation has clear benefits.⁴⁻⁷ For neck pain, however, there are fewer studies and the findings vary, but there is some evidence that conclude spinal manipulation benefits patients presenting with neck pain.⁸⁻¹³ This evidence indicates that the benefits of spinal manipulation include relief of acute neck pain, and reduction in neck pain as measured by validated instruments in sub-acute and chronic neck pain compared with muscle relaxants or usual medical care.¹³⁻¹⁷ Bronfort et al.¹⁵ specifically concluded that for

patients with chronic neck pain, there is moderate evidence that (1) manipulation and mobilization are superior to general practitioner management in the short term, (2) high-technology exercise results in more pain improvement than manipulation in the long term for a mix of patients with acute and chronic pain, and (3) mobilization is superior to physical therapy and general medical care and similar to manipulation in both the short and long term.

Benefits of spinal manipulation for areas beyond the low back and neck include short-term relief from tension-type headaches.¹⁴ Manipulation relieves cervicogenic headache and is comparable to commonly used first line prophylactic prescription medications for tension-type headache and migraine.

Harm

Overall

The literature presents varying conclusions on the harms of spinal manipulative treatment (SMT).^{6,7} In a 2017 review of risks associated with spinal manipulation, 46% percent of the studies reviewed found spinal manipulation to be safe, 42% percent were neutral (did not find harm/benefit); and the remaining 12% percent concluded that spinal manipulation was unsafe because of the possibility of serious adverse events.⁷ Nevertheless, the existence of any adverse effect should not be trivialized.

Studies have noted that there are two types of adverse effects as a result of SMT. The first type is considered to be mild adverse events that are short-term and non-serious such as dizziness, fatigue, and muscle soreness/ discomfort.^{7,18} These side effects occur in 23-83% of patients. The second type of adverse events is more serious and includes cervical artery dissection, stroke, spinal cord injuries, and other serious conditions outcomes related to vertebrobasilar accidents (VBAs). Currently, much of the literature discusses vertebrobasilar insufficiency or vertebralbasilar ischemia (VBI) which is a type of VBA and is often determined to be the link to the more serious adverse events. Nonetheless, serious adverse events are seen as a rarity, and it is estimated that they occur in the range of every 20,000 to 250,000,000 manipulation performed.^{7,18-27}

Most of the reported cases of adverse outcomes have involved thrust or High Velocity/Low Amplitude (HVLA) types of manipulative treatment.^{18,25} Unfortunately, many of the reported cases do not distinguish the type of manipulative treatment provided.

VBAs

VBAs account for 1.3 in 1000 cases of stroke, making them a rare event. Approximately 5% of patients with a VBA die as a result, while 75% have a good functional recovery.²⁸ The most common risk factors for VBAs are migraine, hypertension, oral contraceptive use and smoking.²⁹ Elevated homocysteine levels, which have been implicated in cardiovascular disease, may be a risk factor for a VBA.³⁰

The risk of a VBA occurring spontaneously, is nearly twice the risk of a VBA resulting from cervical spine manipulation.¹⁴ A study done in 1999 reviewing 367 cases of VBA reported from 1966-1993 showed 115 cases related to cervical spine manipulation; 167 were spontaneous, 58 from trivial trauma and 37 from major trauma.³¹

A study in 2002 conducted by Haldeman et al., reported that a VBA following cervical spine manipulation was unpredictable.¹⁴ The authors, however, concluded that a VBA following cervical spine manipulation was "idiosyncratic and rare". Further review of the data showed that 25% of the cases presented with sudden onset of new and unusual headache and neck pain often associated with other neurologic symptoms that may have represented a dissection in progress.³²

Complications from cervical spine manipulation most often occur in patients who have had prior manipulation uneventfully and without obvious risk factors for a VBA.¹⁴ “Most vertebrobasilar artery dissections occur in the absence of cervical manipulation, either spontaneously or after trivial trauma or common daily movements of the neck, such as backing out of the driveway, painting the ceiling, playing tennis, sneezing, or engaging in yoga exercises.”²⁴ In some cases manipulation may not be the primary culprit for causing the dissection, but an aggravating factor or coincidental event.³²

It has been proposed that thrust techniques that use a combination of hyperextension, rotation and traction of the upper cervical spine will place the patient at greatest risk of injuring the vertebral artery. In a retrospective review of 64 medical legal cases, information on the type of manipulation was available in 39 (61%) of the cases. Fifty-one percent (51%) involved rotation, with the remaining 49% representing a variety of positions including lateral flexion, traction and isolated cases of non-force or neutral position thrusts. Only 15% reported any form of extension.³²

Cervical Artery Dissection (CAD)

CAD occurs at a rate of 2.9 per 100,000 individuals every year in the general population, and a large majority (89%) of the individuals diagnosed with CAD have no symptoms or no significant disability that prohibits them from being productive within the following three months of the event.³³ Among those with symptoms, headaches and neck pain are the predominant symptoms for CAD. This creates a dilemma for physicians because cervical spine manipulation is often sought to treat these medical issues. Thus, it is difficult to determine if manipulation causes CAD or if CAD existed at the time of treatment.

Limitations of Studies and Concerns with Pre-manipulation Screening

Due to the design of studies (case reports or retrospective surveys), infrequent reporting of adverse events, and the rare occurrence of many of the more serious complications, it is difficult to determine a causal relationship between SMT and the serious adverse effect.^{7,33} Thus the lingering question of whether or not pre-existing pathologies may have existed prior to the patient receiving SMT remains.^{18,26,34}

In Malone et al., the authors reported that cervical spine manipulation may worsen preexisting cervical disc herniation or even cause cervical disc herniation.²⁶ This report describes complications such as radiculopathy, myelopathy, and vertebral artery compression by a lateral cervical disc herniation. The incidence of these types of complications could be lessened by rigorous adherence to published exclusion criteria for cervical spine manipulation.^{26,35}

Another noteworthy point to highlight is that the literature does not clearly distinguish the type of provider (i.e. M.D., D.O., D.C. or P.T.) or manipulative treatment (manipulation vs. mobilization) provided in cases associated with serious adverse effects. This information may help to understand the mechanism of injury leading to serious adverse effects, as there are differences in education and practice among the various professions that utilize this type of treatment. It is duly noted that the osteopathic approach strictly limits the “thrust”, which is more commonly referred to as “impulse” in osteopathic practicums, to the physiologic barrier as opposed to the chiropractic approach may extend to the parapsysiologic space.

Additionally, pre-manipulation screening tools, that might be used to identify a patient’s risk for VBA and cervical artery dissection have been widely criticized because they have been found to be unreliable and difficult to validate.^{28,29,36-43} These studies have examined the DeKleyn’s test and others like it and determined the tests are unreliable

for demonstrating reproducibility of ischemia or risk of injuring the vertebral artery.³⁶⁻⁴³ For this reason, researchers and groups such as the Bone and Joint Decade Task Force on Neck Pain and Its Associated Disorders recommend that all health care providers conduct a thorough patient history, physical examination and patient self-assessment to rule out certain pre-existing conditions.^{13,44}

Alternative Treatments

Non-steroidal anti-inflammatory drugs (NSAIDs)

NSAIDs such as ibuprofen and aspirin are the most commonly prescribed medications for neck pain. More than 30 million people worldwide use NSAIDs regularly.⁴⁵ In fact, 5% of all medical visit outcomes in the U.S. include a prescription for NSAIDs.⁴⁶ NSAIDs offer temporary relief, but long-term use, gender, age, strength of dose as well as consumption of multiple medications simultaneously may be associated with serious risks affecting the gastrointestinal (GI), renal and cardiovascular systems.^{47,48} Eighty-one percent (81%) of GI bleeds related to NSAID use occur without prior symptoms.⁴⁹ Research in the United Kingdom has shown NSAIDs will cause 12,000 emergency admissions and 2,500 deaths per year due to GI tract complications.³⁰ The annual cost of GI tract complications in the U.S. is estimated at \$3.9 billion, with up to 103,000 hospitalizations and at least 16,500 deaths per year therein making GI toxicity from NSAIDs the 15th most common cause of death in the United States.⁴⁹⁻⁵¹

Epidural steroid injections

Epidural steroid injections (ESIs) are a popular treatment for neck pain.⁵⁰ Complications to ESIs generally occur because of needle placement or drug administration. Common risks associated with needle placement include subdural injection, intrathecal injection and intravascular injection.⁵¹ Subdural injection occurs in ~ 1% of procedures, intrathecal injection occurs in ~ 0.6-10.9% of procedures, and intravascular injection, the most significant risk, occurs in ~ 2% of procedures.⁵¹ Other risks include cervical epidural abscess, dural puncture, spinal cord trauma, infection, hematoma, nerve damage, vascular injury and cerebral vascular or pulmonary embolus.^{52,53} Complications that may arise from drug administration include osteoporosis, Cushing's syndrome, avascular necrosis of bone, and steroid myopathy. While complications due to needle placement or administration of steroids are rare, they have been reported in the literature.^{52,53}

Conclusion

Osteopathic manipulative treatment of the cervical spine, including but not limited to HVLA treatment, is effective for low back and neck pain and is safe. Because of the rarity of serious adverse events, trainees and practicing physicians should be provided with sufficient information so they are advised of the potential risks and able to communicate the potential risks to their patients. Prior to recommending cervical spine manipulations, physicians should conduct a thorough patient exam and medical history review to try to identify any preexisting conditions that may indicate the patient is at risk for a serious adverse event. Additionally, it is recognized that there is a need for research to distinguish the risk of VBA and CAD associated with manipulation done by specific provider types as well as research to determine the nature of the relationship between the different types of manipulative treatment and VBA and CAD.

It is the position of the American Osteopathic Association that all modalities of osteopathic manipulative treatment of the cervical spine, including HVLA, should continue to be taught at all levels of education, and that osteopathic physicians should continue to offer this form of treatment to their patients. Physicians should use a combination of medical history reviews and physical exams, diagnostic studies, and best

judgment to determine if a patient has any pre-existing conditions that place the patient at risk of suffering a serious adverse event.

Special Acknowledgements

In crafting the updated Position Statement, the Bureau of Osteopathic Clinical Education and Research (BOCER) would like to thank the Osteopathic Manipulation Medicine and Osteopathic Manipulative Treatment (OMM/OMT) Research Task Force for its input, and a special thank you to Hollis King, DO, PhD, who served as an outside contributor.

References

1. Gaskin D.J., and Richard, P. (2012). The economic costs of pain in the United States. *The Journal of Pain*, 13(8), 715-724.
2. US Burden of Disease Collaborators. (2013). The state of US health, 1990–2010 burden of diseases, injuries, and risk factors. *Journal of American Medical Association*, 310(6), 591–608.
3. GBD. (2017). Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*, 390, 1211-1259.
4. Shekelle, P, Adams, A, et al. (1992). Spinal manipulation for low-back pain. *Annals of Internal Medicine*, 117(7), 590-98.
5. Rubinstein, S.M., de Zoete, A., van Middelkoop, M., Assendelft, W.J.J., de Boer, M.R., and van Tulder, M.W. (2019). Benefits and harms of spinal manipulative therapy for the treatment of chronic low back pain: Systematic review and meta-analysis of randomized controlled trials. *BMJ*, 364, 1689, <https://doi.org/10.1136/bmj.l689>.
6. Paige, N., Miake-Lye, I.M., Booth, M.S., Beroes, J.M., Mardian, A.S., Dougherty, P., Branson, R., Tang, B., Morton, S.C., and Shekelle, P.G. (2017). Association of spinal manipulative therapy with clinical benefit and harm for acute low back pain systematic review and meta-analysis. *Journal of American Medical Association*, 317(14), 1451-1460.
7. Nielsen, S.M., Tarp, S. Christensen, R., Bliddal, H., Klokke, L., and Henriksen, M. (2017). The risk associated with spinal manipulation: An overview of reviews. *Systematic Reviews*, 6, 64-83.
8. Koes, B.W., Bouter, L.M., van Marmeren, H., Essers, A.H., Verstegen, G.M., Hofhuizen, D.M., Houben, J.P., and Knipschild, P.G. (1992). The effectiveness of manual therapy, physiotherapy, and treatment by the general practitioner for nonspecific back and neck complaints, a randomized clinical trial. *Spine*, 17(1), 28-35.
9. Koes, B.W., Bouter, L.M., van Marmeren, H., Essers, A.H., Verstegen, G.M., Hofhuizen, D.M., Houben, J.P., and Knipschild, P.G. (1992). Randomised clinical trial of manipulative therapy and physiotherapy for persistent back and neck complaints: Results of one year follow up. *BMJ*, 304, 601-605.
10. Koes, B.W., Bouter, L.M., van Marmeren, H., Essers, A.H., Verstegen, G.M., Hofhuizen, D.M., Houben, J.P., and Knipschild, P.G. (1993). A randomized clinical trial of manual therapy and physiotherapy for persistent neck and back complaints: sub-group analysis and relationship between outcome measures. *Journal of Manipulative and Physiological Therapeutics*, 16(4), 211-219.
11. Cassidy, J.D., Lopes, A.A., and Yong-Hing, K. (1992). The immediate effect of manipulation versus mobilization on pain and range of motion in the cervical spine: A

- randomized controlled trial. *Journal of Manipulative and Physiological Therapeutics*, 15, 570-575.
12. Jensen, O.K., Nielsen, F.F., and Vosmar, L. (1990). An open study comparing manual therapy with the use of cold packs in the treatment of posttraumatic headache. *Cephalgia*, 10, 241-250.
 13. Hadleman, S., Carroll, L., Cassidy, J.D., Schubert, J., and Nygren, A. (2008). The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine*, 33 (4S), S5-S7.
 14. Hurwitz, E.L., Aker, P.D., Adams, A.H., Meeker, W.C., and Shekelle, P.G. (1996). Manipulation and mobilization of the cervical spine. A systematic review of the literature. *Spine*, 21(15), 1746-56.
 15. Bronfort, G., Haas, M, Evans, R.L., and Bouter, L.M. (2004). Efficacy of spinal manipulation and mobilization for low back pain and neck pain: A systematic review and best evidence synthesis. *Spine Journal*, 4, 335-356.
 16. Garcia, J.D., Arnold, S., Tetley, K., Voight, K., and Frank, R.A. (2016). Mobilization and manipulation of the cervical spine in patients with cervicogenic headache: Any scientific evidence? *Frontiers in Neurology*, 7 (40).
 17. Gross, A.R., Aker, P.D., Goldsmith, C.H., and Peloso, P. (1996). Conservative management of mechanical neck disorders: A systematic overview and meta-analysis. *BMJ*, 313 (7068), 1291-1296.
 18. Swait, G. and Finch, R. (2017). What are the risks of manual treatment of the spine? A scoping review for clinicians. *Chiropractic and Manual Therapies*, 25, 37-52.
 19. Jensen, O.K., Nielsen, F.F., and Vosmar, L. (1990). An open study comparing manual therapy with the use of cold packs in the treatment of posttraumatic headache. *Cephalgia*, 10, 241-250.
 20. Koss, R.W. (1990). Quality assurance monitoring of osteopathic manipulative treatment. *Journal of the American Osteopathic Association*, 90(5), 427-433.
 21. Dvorak, J., and Orelli, F. (1985). How dangerous is manipulation to the cervical spine? Case report and results of a survey. *Manual Medicine*, 2, 1-4.
 22. Carey P.F. (1993). A report on the occurrence of cerebral vascular accidents in chiropractic practice. *The Journal of the Canadian Chiropractic Association*, 37,104-106.
 23. Coulter, I.D., Hurwitz, E., Adams, A.H., Meeker, W., Hansen, D.T., Mootz, R., Aker, P., Genovese, B., and Shekelle, P.G (1996). The appropriateness of manipulation and mobilization of the cervical spine. Santa Monica, CA: RAND Corporation, https://www.rand.org/pubs/monograph_reports/MR781.html.
 24. Haldeman, S., Kohlbeck, F.J., and McGregor, M. (2002). Unpredictability of cerebrovascular ischemia associated with cervical spine manipulation: A review of 64 cases after cervical spine manipulation therapy. *Spine*, 27, 49-55.
 25. Assendelft, W.J.J., Bouter, L.M., and Knipschild, P.G. (1996). Complications of spinal manipulation: A comprehensive review of the literature. *Journal of Family Practice*, 42, 475-480.
 26. Malone, D.G., Baldwin, N.G., Tomecek, F.J., Boxell, C.M., Gaede, S.E., Covington, C.G., and Kugler, K.K. (2002). Complications of cervical spine manipulation therapy: 5-Year retrospective study in a single-group practice. *Journal of Neurosurgery*, DOI: <https://doi.org/10.3171/foc.2002.13.6.8>.
 27. Vick, D.A., McKay, C., and Zengerle, C.R. (1996). The safety of manipulative treatment: Review of the literature from 1925 to 1993. *Journal of American Osteopathic Association*, 96(2), 113-115.

28. Schievink, W. (2001). Spontaneous dissection of the carotid and vertebral arteries. *New England Journal of Medicine*, 344(12), 898-906.
29. Blower, A.I., Brooks, A., Fenn, C.G., and Hill, A. (1997). Emergency admissions for upper gastrointestinal disease and their relation to NSAID use. *Alimentary Pharmacology and Therapeutics*, 11(2), 283-291.
30. Rosner, A. (2004). Spontaneous Cervical artery dissections and implications for homocysteine. *Journal of Manipulative and Physiological Therapeutics*, 27(2), 124-132. (35)
31. Haldeman, S., Kohlbeck, F.J., and McGregor, M. (1999). Risk factors and precipitating neck movements causing vertebrbasilar artery dissection after cervical trauma and spinal manipulation. *Spine*, 24, 785-794.
32. Haldeman, S., Kohlbeck, F.J., and McGregor, M. (2002). Stroke, cerebral artery dissection, and cervical spine manipulative therapy. *Journal of Neurology*, 249(8), 1098-1104.
33. Chaibi, A., and Russell, M.B. (2019). A risk-benefit assessment strategy to exclude cervical artery dissection in spinal manual-therapy: A comprehensive review. *Annals of Medicine*, DOI: 10.1080/07853890.2019.1590627.
34. Schwartz, N.E., Vertinsky, A.T., Hirsch, K.G., and Albers, G.W. (2009). Clinical and radiographic natural history of cervical artery dissections. *Journal of Stroke and Cerebrovascular Disease*, 18(6), 416-423.
35. Childs, J.D., Flynn, T.W., Fritz, J.M., Piva, S.R., Whitman, J.M., Wainner, R.S., and Greenman, P.E. (2005). Screening for vertebrbasilar insufficiency in patients with neck pain: Manual therapy decision-making in the presence of uncertainty. *Journal of Orthopaedic and Sports Physical Therapy*, 35, 300-306.
36. Licht, P.B., Christensen, H.W., Svendsen, P., and Hoilund-Carlsen, P.F. (1999). Vertebral artery flow and cervical manipulation: An experimental study. *Journal of Manipulative and Physiological Therapeutics*, 22 (7), 431-435.
37. Cote, P., Kreitz, B.G., Cassidy, J.D., and Theil, H. (1996). The validity of extension-rotation tests as a clinical screening procedure before neck manipulation: A secondary analysis. *Journal of Manipulative and Physiological Therapeutics*, 19, 159-164.
38. Refshauge, K.M. (1994). Rotation: A valid premanipulative dizziness test? Does it predict safe manipulation? *Journal of Manipulative and Physiological Therapeutics*, 17(1), 15-19.
39. Stevens, A. (1991). A functional doppler sonography of the vertebral artery and some considerations about manual techniques. *Journal of Manual Medicine*, 6, 102-105.
40. Theil, H., Wallace, K., Donat, J., Yong-Hing, K. (1994). Effect of various head and neck positions on vertebral artery blood flow. *Clinical Biomechanics*, 9, 105-110.
41. Weingart, J.R., and Bischoff, H.P. (1992). Doppler sonography of the vertebral artery with regard to head positions appropriate to manual medicine. *Manuelle Medizin*, 30, 62-65.
42. Rivett, D.A., Sharples, K.J. and Milburn, P.D. (1999). Effect of premanipulative tests on vertebral artery and internal carotid artery blood flow: A pilot study. *Journal of Manipulative and Physiological Therapeutics*, 22 (6), 368-375.
43. Kerry, R., Taylor, A.J., Mitchell, J., McCarthy, C., and Brew, J. (2008). Manual therapy and cervical arterial dysfunction, directions for the future: A clinical perspective. *The Journal of Manual and Manipulative Therapy*, 16 (1), 39-48.
44. Puentedura, E.J., Cleland, J.A., Landers, M.R., Mintken, P., Louw, A., and Fernandez-de-las-Penas, C. (2012). Development of a clinical prediction rule to identify patients with

- neck pain likely to benefit from thrust joint manipulation to the cervical spine. *Journal of Orthopaedic and Sports Physical Therapy*, 42(7), 577-591.
45. Gargiulo, G., Capodanno, D., Longo, G., Capranzano, P., and Tamburino, C. (2014). Updates on NSAIDs in patients with and without coronary artery disease: Pitfalls, interactions and cardiovascular outcomes. *Expert Review of Cardiovascular Therapy*, 12 (10), 1185-1203.
 46. Gunter, B.R., Butler, K.A., Wallace, R.L., Smith, S.M., and Harirforoosh, S. (2017). Non-steroidal anti-inflammatory drug-induced cardiovascular adverse events: A meta-analysis. *Journal of Clinical Pharmacy and Therapeutics*, 42, 27-38.
 47. Shep, D., Khanwelkar, C., Gade, P., and Larad, S. (2019). Safety and efficacy of curcumin versus diclofenac in knee osteoarthritis: A randomized open-label parallel-arm study. *Trials*, 20 (1), 214-225.
 48. Moore, N., Pollack, C., and Butkerait, P. (2015). Adverse drug reactions and drug-drug interactions with over-the-counter NSAIDs. *Therapeutics and Clinical Risk Management*, 11, 1061-1075.
 49. Wolfe, M., Lichtenstein, D., and Singh, G. (1999). Gastrointestinal toxicity of nonsteroidal anti-inflammatory drugs. *New England Journal of Medicine*, 340(24), 1888-1899.
 50. McGrath, J.M., Schaefer, M.P., and Malkamaki, D.M. (2011). Incidence and characteristics of complications from epidural steroid injections. *Pain Medicine*, 12, 726-731.
 51. Mulroy, M., Norris, M., and Spencer L. (1997). Safety steps for epidural injection of local anesthetics: Review of the literature and recommendations. *Anesthesia & Analgesia*, 85(6), 1346-1356.
 52. Huang, R.C., Shapiro, G.S., Lim, M., Sandhu, H.S., Lutz, G.E., and Herzog, R.J., (2004). Cervical epidural abscess after epidural steroid injection. *Spine*, 29(1), E7-E9.
 53. Abdi, S., Datta, S., Trescot, A.M., Schultz, D.M., Adlaka, R., Alturi, S.L., Smith, H.S., and Manchikanti, L. (2007). Epidural steroids in the management of chronic spinal pain: A systematic review. *Pain Physician*, 10, 185-212.

Source: H324-A/19

Status: 2004; 2009 Reaffirmed; 2014 Reaffirmed; 2019 Reaffirmed